

OBSTETRICS

Maternal and newborn outcomes in planned home birth vs planned hospital births: a metaanalysis

Joseph R. Wax, MD; F. Lee Lucas, PhD; Maryanne Lamont, MLS; Michael G. Pinette, MD; Angelina Cartin; Jacquelyn Blackstone, DO

OBJECTIVE: We sought to systematically review the medical literature on the maternal and newborn safety of planned home vs planned hospital birth.

STUDY DESIGN: We included English-language peer-reviewed publications from developed Western nations reporting maternal and newborn outcomes by planned delivery location. Outcomes' summary odds ratios with 95% confidence intervals were calculated.

RESULTS: Planned home births were associated with fewer maternal interventions including epidural analgesia, electronic fetal heart rate monitoring, episiotomy, and operative delivery. These women were less

likely to experience lacerations, hemorrhage, and infections. Neonatal outcomes of planned home births revealed less frequent prematurity, low birthweight, and assisted newborn ventilation. Although planned home and hospital births exhibited similar perinatal mortality rates, planned home births were associated with significantly elevated neonatal mortality rates.

CONCLUSION: Less medical intervention during planned home birth is associated with a tripling of the neonatal mortality rate.

Key words: neonatal mortality, patient safety, planned home childbirth

Cite this article as: Wax JR, Lucas FL, Lamont M, et al. Maternal and newborn outcomes in planned home birth vs planned hospital births: a metaanalysis. *Am J Obstet Gynecol* 2010;203:x-ex-x.ex.

Approximately 1 in 200 US women deliver at home, accounting for approximately 25,000 deliveries annually.¹ An estimated 75% of low-risk singleton home births appear to be planned home deliveries.² The American College of Obstetricians and Gynecologists does not support home birth, citing safety concerns and lack of rigorous scientific study.³ Ideally, further investigation regarding the relative

From the Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology (Drs Wax, Pinette, and Blackstone and Ms Cartin); the Maine Center for Outcomes Research and Evaluation (Dr Lucas); and the Medical Library (Ms Lamont), Maine Medical Center, Portland, ME.

Presented at the 30th Annual Meeting of the Society for Maternal-Fetal Medicine, Chicago, IL, Feb. 1-6, 2010.

Received Jan. 20, 2010; revised April 14, 2010; accepted May 18, 2010.

Reprints not available from the authors.

0002-9378/free

© 2010 Mosby, Inc. All rights reserved.

doi: 10.1016/j.ajog.2010.05.028



For Editors' Commentary,
see Table of Contents

★ EDITORS' CHOICE ★

safety of planned home vs planned hospital delivery would occur via randomized trials, which are, however, impractical. Large cohort studies comparing outcomes of actual home with actual hospital births provide valuable data, particularly regarding rare but serious events.² However, such investigations likely underestimate the risks associated with planned home birth, as up to 9% of parous and 37% of nulliparous women intending home birth require in-trapartum transfer to hospital.⁴⁻⁷ Thus, adverse outcomes among the latter deliveries are attributed to hospital births. Therefore, cohort studies comparing planned home with planned hospital births provide the only sources of data by intended delivery location. Since individual reports of this design are limited by sample size, we employed metaanalysis according to proposed reporting methods to clarify the relative merits of planned home vs planned hospital birth.⁸

MATERIALS AND METHODS

Search strategy

Computerized literature searches of MEDLINE and EMBASE were performed by a physician and medical librarian.

MEDLINE search results

The search strategy for the query for "all studies, regardless of methods, comparing intended/planned home births to intended/planned hospital births for maternal and newborn outcomes" was run in the MEDLINE database from 1950 through November week 1 2009 (Figure 1). The following terms were used: explosion of the medical subject heading "Home Childbirth" (defined as childbirth taking place at home); explosion of the medical subject heading "Delivery, Obstetric" (defined as delivery of the fetus and placenta under the care of an obstetrician or a health worker; obstetric deliveries may involve physical, psychological, medical, or surgical interventions); explosion of the medical subject heading "Hospitalization" (defined as being in a hospital or being placed in a hospital); the confinement of a patient in a hospital); and explosion of the medical subject heading "Inpatients" (defined as persons admitted to health facilities that provide board and room, for the purpose of observation, care, diagnosis, or treatment). The terms "Hospitalization" or "Inpatients" or any mention of the word form "Hospital*" (designated with an asterisk as the wild card picking up any

FIGURE 1
MEDLINE search strategy

Database: Ovid MEDLINE(R) <1950 to November Week 1 2009>
Search Strategy:

- ```

1 exp Home Childbirth/ (1495)
2 exp Delivery, Obstetric/ (54357)
3 exp Hospitalization/ (122627)
4 exp Inpatients/ (8376)
5 2 and (3 or 4 or hospital*.mp) [mp=title, original title, abstract, name of substance
word, subject heading word, unique identifier] (7942)
6 1 and 5 (155)
7 6 (155)
8 limit 7 to (english language and humans) (132)
9 8 and (outcome* or compar* or intend* or plan*).mp. [mp=title, original title,
abstract, name of substance word, subject heading word, unique identifier] (83)
10 from 9 keep 1-83 (83)

```

Wax. *Outcomes in planned home birth vs planned hospital births. Am J Obstet Gynecol* 2010.

letters after the “l,” eg, “hospitals,” “hospitalized”) was then combined with the term “Delivery, Obstetric” to limit to a hospital birth. These results were then “anded” with the term “Home Childbirth” and by doing so indicated that the citation must include indexing for both terms; thus the discussion in the article would include both concepts. Limits to English language and human studies were then included. The final line of strategy was to take the retrieval and limit to any citations that would include the word forms for “outcome\*” or “compar\*” or “intend\*” or “plan\*” as a way to narrow the results to include the concepts of outcomes, comparisons, comparing, intended, or planned by using the asterisk as a wild card.

**EMBASE search results**

This strategy was done using EMBASE classic (1947 through present). Using the all subject words feature the term “Home Delivery” was searched. The term “Childbirth” was also searched and combined with any form of the word “Hospital?” with the ? indicating a wild card to pick up any forms of the word, such as “hospitals” and “hospitalization.”

The Cochrane Database of Systematic Reviews was also searched for relevant publications. Titles and abstracts of citations were reviewed for potential relevance and selected manuscripts were reviewed. References in these papers were

manually reviewed and retrieved if potentially relevant.

**Study selection criteria**

Inclusion criteria were determined before the literature search was performed. Studies were included if performed in developed Western countries, published in English-language peer-reviewed literature, maternal and newborn outcomes were analyzed by planned delivery location, and data were presentable in a 2×2 table. Manuscripts were evaluated for quality using a published instrument.<sup>9</sup> Outcome data were extracted by 2 physicians, with differences resolved by consensus. Outcomes for maternal intervention included epidural analgesia, electronic fetal heart rate monitoring, episiotomy, operative vaginal delivery (forceps or vacuum), and cesarean delivery. Maternal outcomes included mortality, morbidity measures of lacerations (≥3 degrees, vaginal, and perineal), infections (chorioamnionitis, endometritis, wound, and urinary), postpartum hemorrhage, retained placenta, and umbilical cord prolapse. Neonatal outcomes included 5-minute Apgar score <7, prematurity (<37 weeks’ gestation), low birthweight (<10% for gestational age or <2500 g), macrosomia (≥90% for gestational age or ≥4000 g), postdatism (≥42 weeks’ gestation), assisted ventilation requirement, perinatal death (stillbirth of at least 20 weeks or 500 g or

death of liveborn within 28 days of birth), and neonatal death (death of a liveborn within 28 days of delivery). Perinatal and neonatal deaths were evaluated overall and for nonanomalous offspring. The study did not require institutional review board approval.

**Statistical methods**

Studies were assessed for homogeneity using the Breslow-Day test. When present, a fixed effects model was used; when absent, a random effects model was employed. Summary odds ratios (ORs) with 95% confidence intervals (CIs) were calculated for maternal and newborn outcomes, comparing planned home to planned hospital deliveries. Sensitivity analyses were conducted for studies employing matched planned home and hospital births,<sup>4,10-12</sup> those primarily based upon pre-1990 data,<sup>5,10,13</sup> lesser quality reports,<sup>5,13,14</sup> and those not clearly specifying home birth attendants or in which home births were conducted by other than certified or certified nurse midwives.<sup>10,15</sup> We used software (SAS, version 9.2; SAS Institute Inc, Cary, NC) for most data analysis. Random effects results were analyzed using an online metaanalysis calculator from the University of Pittsburgh (<http://www.pitt.edu/~super1/lecture/lec1171/meta5.doc>).

**RESULTS**


The results of the literature search are noted in Figure 2. Characteristics of the 12 included studies are described in Table 1.<sup>4-7,10-17</sup> A total of 342,056 planned home and 207,551 planned hospital deliveries were available for analysis. No maternal deaths were reported in 4 studies totaling 10,977 planned home and 28,501 planned hospital births, precluding metaanalysis. However, we calculated the upper 95% confidence limits for these rates, expressed per 100,000 births, as 27.3 and 10.5, respectively.<sup>4,7,11,12</sup> Table 2 presents the metaanalysis of maternal outcomes by intended delivery location. Planned home births experienced significantly fewer medical interventions including epidural analgesia, electronic fetal heart rate monitoring, episiotomy, and operative vaginal and cesarean deliveries. Likewise,

women intending home deliveries had fewer infections,  $\geq 3$ -degree lacerations, perineal and vaginal lacerations, hemorrhages, and retained placentas. There was no significant difference in the rate of umbilical cord prolapse.

Table 3 describes the metaanalysis of neonatal outcomes. Low Apgar scores could not be evaluated as most studies considered thresholds other than a score of 7 (range, 4–8). Compared to offspring of women planning hospital births, those of mothers planning home births were less likely to be born preterm or be of low birthweight. However, planned home births more often progressed to  $\geq 42$  weeks. While there was no difference in the rate of assisted ventilation, 1 large study found more frequent ventilation among planned home births, while 2 smaller studies noted lower rates in this group.<sup>11,15,17</sup> Perinatal mortality was similar by intended delivery location, overall as well as just among nonanomalous offspring. In contrast, the overall neonatal death rate was almost twice as high in planned home vs planned hospital births, and almost tripled among nonanomalous neonates. Importantly, these latter observations were consistent across all studies examining neonatal mortality, regardless of the covered time period.<sup>4,7,10,13,15,17</sup> The anticipated population-based attributable risk of neonatal death overall and among nonanomalous offspring, employing a home birth prevalence of 0.6%, was 0.3% and 0.4%, respectively.

The results of the sensitivity analyses excluding older studies and poorer quality investigations revealed no significantly different findings from the original metaanalysis. In contrast, the sensitivity analysis excluding the 4 papers employing matching found no significant differences between planned home and planned hospital births regarding  $\geq 3$ -degree lacerations (OR, 0.90; 95% CI, 0.62–1.31), retained placentas (OR, 0.66; 95% CI, 0.38–1.14), hemorrhage (OR, 0.80; 95% CI, 0.64–1.00), prematurity (OR, 0.52; 95% CI, 0.27–1.00), and neonatal death among nonanomalous offspring (OR, 2.22; 95% CI, 0.83–5.97). The analysis excluding studies that included home births at-

**FIGURE 2**  
Study selection process



Wax. Outcomes in planned home birth vs planned hospital births. *Am J Obstet Gynecol* 2010.

tended by other than certified or certified nurse midwives had findings similar to the original study, except that the ORs for neonatal deaths among all (OR, 1.57; 95% CI, 0.62–3.98) and nonanomalous (OR, 3.00; 95% CI, 0.61–14.88) newborns were not statistically significant.

### COMMENT

Of concern, this investigation identified a doubling and tripling of the neonatal mortality rate overall and among nonanomalous offspring, respectively, in planned home compared to planned hospital births. This finding is particularly robust considering the homogeneity of the observation across studies. It is especially striking as women planning

home births were of similar and often lower obstetric risk than those planning hospital births. The planned home delivery group commonly exhibited fewer obstetric risk factors such as excessive body mass index, nulliparity, prior cesarean, and previous pregnancy complications.<sup>6,7,10,17</sup> Moreover, our data show that planned home births are characterized by less frequent premature and low birthweight infants. The differential obstetric risk by planned delivery location was not unexpected since women self-select for home birth.

In developed nations, following congenital anomalies, most perinatal deaths are related to intrapartum anoxia.<sup>18</sup> Among the studies in our meta-

**TABLE 1**  
**Characteristics of studies included in metaanalysis**

| Setting                                 | Study design                          | Time period studied | Publication year | Data source                                                                      | Inclusions                                                                                                                                                                                                  | Data analysis by parity | Planned deliveries, n |          | Intrapartum transfer to hospital rate |             |              |
|-----------------------------------------|---------------------------------------|---------------------|------------------|----------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------|-----------------------|----------|---------------------------------------|-------------|--------------|
|                                         |                                       |                     |                  |                                                                                  |                                                                                                                                                                                                             |                         | Home                  | Hospital | Nulliparous                           | Parous      | Overall      |
| California, United States <sup>13</sup> | Retrospective cohort                  | 1976-1982           | 1984             | ND                                                                               | Single obstetrician and lay midwife practice, nulliparous and parous                                                                                                                                        | Stratification          | 454                   | 67       | 25/258 (9.7)                          | 5/263 (1.9) | 30/521 (5.8) |
| United Kingdom <sup>5</sup>             | Prospective cohort                    | 1978-1983           | 1985             | Submitted data collection forms                                                  | Low risk, parous, no past obstetric complications, 26 practices                                                                                                                                             | Parous only             | 202                   | 185      | ND                                    | 3.5%        | 3.5%         |
| Western Australia <sup>10</sup>         | Matched cohort                        | 1981-1987           | 1994             | Birth records, transfer forms, computer system                                   | All Western Australian women booking for home birth and matched cohort of not planned home birth, nulliparous and parous                                                                                    | Matching                | 976                   | 2928     | ND                                    | ND          | 14.0%        |
| Switzerland <sup>4</sup>                | Prospective cohort with matched pairs | 1989-1992           | 1996             | Special data collection forms                                                    | Women receiving care from 1 team of physicians and midwives, no formal policy for planned home delivery, nulliparous and parous                                                                             | Matching                | 489                   | 385      | 25%                                   | ND          | 15.9%        |
| Netherlands <sup>5</sup>                | Prospective cohort                    | 1990-1993           | 1996             | Questionnaire, birth records                                                     | Low-risk pregnancies receiving midwifery care in 54 practices, nulliparous and parous                                                                                                                       | Stratification          | 1140                  | 696      | 36.7%                                 | 8.7%        | 20.3%        |
| Sweden <sup>7</sup>                     | Population-based cohort               | 1992-2004           | 2008             | Swedish Medical Birth Register                                                   | All Swedish women planning home birth and control group of 37-42 wk low-risk singletons in ratio of 1:10, nulliparous and parous                                                                            | No                      | 897                   | 11,341   | ND                                    | ND          | ND           |
| British Columbia, Canada <sup>17</sup>  | Prospective cohort                    | 1998-1999           | 2002             | British Columbia Reproductive Care Program antenatal, birth, and newborn records | Low-risk women $\geq 36$ wk planning home birth with midwife enrolled in Home Birth Demonstration Project and low-risk women 37-41 wk planning hospital birth, physician or midwife, nulliparous and parous | No                      | 862                   | 1314     | ND                                    | ND          | 16.5%        |
| United Kingdom <sup>14</sup>            | Randomized trial                      | 1994                | 1996             | ND                                                                               | Low-risk parous women in 1 practice                                                                                                                                                                         | Parous only             | 5                     | 6        | 0                                     | 0           | 0            |

Wax. Outcomes in planned home birth vs planned hospital births. *Am J Obstet Gynecol* 2010.

(continued)

**TABLE 1**  
**Characteristics of studies included in metaanalysis** (continued)

| Setting                                       | Study design                                  | Time period studied | Publication year | Data source                           | Inclusions                                                                | Data analysis by parity | Planned deliveries, n |          | Intrapartum transfer to hospital rate |        |         |
|-----------------------------------------------|-----------------------------------------------|---------------------|------------------|---------------------------------------|---------------------------------------------------------------------------|-------------------------|-----------------------|----------|---------------------------------------|--------|---------|
|                                               |                                               |                     |                  |                                       |                                                                           |                         | Home                  | Hospital | Nulliparous                           | Parous | Overall |
| Washington State, United States <sup>15</sup> | Population-based cohort                       | 1989-1996           | 2002             | Birth certificates                    | Low-risk singletons $\geq 34$ wk and $\geq 37$ wk, nulliparous and parous | Adjustment              | 6133                  | 10,593   | ND                                    | ND     | ND      |
| Netherlands <sup>16</sup>                     | Population-based cohort                       | 2000-2006           | 2009             | National perinatal registration data  | Low-risk singletons 37-42 wk, nulliparous and parous                      | Stratification          | 321,307               | 163,261  | ND                                    | ND     | ND      |
| Ontario, Canada <sup>11</sup>                 | Population-based cohort with matched controls | 2003-2006           | 2009             | Ministry of Health midwifery database | Low-risk singletons 37-43 wk, nulliparous and parous                      | Matching                | 6692                  | 6692     | ND                                    | ND     | 5.4%    |
| British Columbia, Canada <sup>12</sup>        | Population-based cohort with matched controls | 2000-2004           | 2009             | Provincial perinatal database         | Low-risk singletons 36-41 wk, nulliparous and parous                      | Not performed           | 2899                  | 10,083   | ND                                    | ND     | ND      |

ND, not described.

Wax. Outcomes in planned home birth vs planned hospital births. *Am J Obstet Gynecol* 2010.

analysis reporting causes of neonatal deaths in planned hospital births, this pattern was confirmed.<sup>4,7,10,15</sup> In contrast, 2 cohort studies implicated intrapartum asphyxia in 31% and 52% of planned home delivery perinatal deaths.<sup>19,20</sup> The past 2 decades have

seen a significant decrease in such deaths, with evidence suggesting fewer fetuses experiencing intrapartum anoxia.<sup>18,21</sup> Speculative explanations for the trend include more liberal use of ultrasound, electronic fetal heart rate monitoring, fetal acid-base assess-

ment, labor induction, and cesarean delivery.<sup>18,21</sup> Our findings, considered in light of these observations, raise the question of a link between the increased neonatal mortality among planned home births and the decreased obstetric intervention in this group.

**TABLE 2**  
**Metaanalysis of maternal outcomes in planned home vs planned hospital births**

| Outcome                                     | No. of studies | Planned home n/N (%) | Planned hospital n/N (%) | OR   | 95% CI    |
|---------------------------------------------|----------------|----------------------|--------------------------|------|-----------|
| <b>Intervention</b>                         |                |                      |                          |      |           |
| Epidural in labor <sup>a</sup>              | 3              | 945/10,453 (9.0)     | 4148/18,089 (22.9)       | 0.24 | 0.22-0.25 |
| Electronic fetal heart rate monitoring      | 2              | 521/3761 (13.8)      | 7138/11,397 (62.6)       | 0.10 | 0.09-0.10 |
| Episiotomy <sup>a</sup>                     | 8              | 939/13,427 (7.0)     | 3075/29,677 (10.4)       | 0.26 | 0.24-0.28 |
| Operative vaginal delivery <sup>a</sup>     | 8              | 497/14,157 (3.5)     | 3433/33,624 (10.2)       | 0.26 | 0.24-0.28 |
| Cesarean delivery <sup>a</sup>              | 10             | 731/14,616 (5.0)     | 3140/33,697 (9.3)        | 0.42 | 0.39-0.45 |
| <b>Morbidity</b>                            |                |                      |                          |      |           |
| $\geq 3$ -degree laceration <sup>a</sup>    | 5              | 150/12,604 (1.2)     | 794/31,740 (2.5)         | 0.38 | 0.33-0.45 |
| Infection <sup>b</sup>                      | 5              | 36/5341 (0.7)        | 319/12,347 (2.6)         | 0.27 | 0.19-0.39 |
| Postpartum bleeding/hemorrhage <sup>a</sup> | 7              | 933/18,720 (4.9)     | 1639/32,552 (5.0)        | 0.66 | 0.61-0.71 |
| Perineal laceration <sup>a</sup>            | 6              | 2408/5632 (42.7)     | 8422/22,695 (37.1)       | 0.76 | 0.72-0.81 |
| Vaginal laceration <sup>b</sup>             | 3              | 640/8078 (7.9)       | 4126/18,418 (22.4)       | 0.85 | 0.78-0.93 |
| Cord prolapse <sup>b</sup>                  | 3              | 3/4658 (0.06)        | 32/22,738 (0.14)         | 0.37 | 0.11-1.24 |
| Retained placenta <sup>a</sup>              | 5              | 73/6079 (1.2)        | 248/15,208 (1.6)         | 0.65 | 0.51-0.83 |

CI, confidence interval; OR, odds ratio.

<sup>a</sup> Random effects model; <sup>b</sup> Fixed effects model.

Wax. Outcomes in planned home birth vs planned hospital births. *Am J Obstet Gynecol* 2010.

TABLE 3

**Metaanalysis of neonatal outcomes in planned home vs planned hospital births**

| Outcome                                               | No. of studies | Planned home n/N (%) | Planned hospital n/N (%) | OR   | 95% CI    |
|-------------------------------------------------------|----------------|----------------------|--------------------------|------|-----------|
| <b>Morbidity</b>                                      |                |                      |                          |      |           |
| Prematurity <37 wk <sup>a</sup>                       | 5              | 75/9751 (0.77)       | 191/4076 (4.7)           | 0.72 | 0.55–0.96 |
| Postdates ≥42 wk <sup>a</sup>                         | 4              | 193/9297 (2.1)       | 238/10,701 (2.2)         | 1.87 | 1.50–2.32 |
| Low birthweight <10% or <2500 g <sup>b</sup>          | 5              | 209/15,411 (1.3)     | 468/21,290 (2.2)         | 0.60 | 0.50–0.71 |
| Large for gestational age >90% or 4000 g <sup>a</sup> | 4              | 1344/13,525 (9.9)    | 1340/17,411 (7.7)        | 1.07 | 0.99–1.16 |
| Newborn ventilation <sup>a</sup>                      | 3              | 497/13,525 (3.7)     | 502/10,701 (4.7)         | 1.12 | 0.99–1.28 |
| <b>Mortality</b>                                      |                |                      |                          |      |           |
| <b>Perinatal death</b>                                |                |                      |                          |      |           |
| All <sup>b</sup>                                      | 6              | 229/331,666 (0.07)   | 140/175,443 (0.08)       | 0.95 | 0.77–1.18 |
| Nonanomalous <sup>b</sup>                             | 4              | 225/330,324 (0.07)   | 134/173,266 (0.08)       | 0.95 | 0.76–1.18 |
| <b>Neonatal death</b>                                 |                |                      |                          |      |           |
| All <sup>b</sup>                                      | 7              | 32/16,500 (0.20)     | 32/33,302 (0.09)         | 1.98 | 1.19–3.28 |
| Nonanomalous <sup>b</sup>                             | 6              | 23/15,633 (0.15)     | 14/31,999 (0.04)         | 2.87 | 1.32–6.25 |

CI, confidence interval; OR, odds ratio.

<sup>a</sup> Random effects model; <sup>b</sup> Fixed effects model.

Wax. Outcomes in planned home birth vs planned hospital births. *Am J Obstet Gynecol* 2010.

Additionally, while limited by the number of neonatal deaths described in sufficient detail, planned home births were characterized by a greater proportion of deaths attributed to respiratory distress and failed resuscitation.<sup>7,10,13,15</sup> These findings echo concerns raised in a recent large US cohort study in which home births experienced significantly more 5-minute Apgar scores <7 as compared to low-risk term hospital births, suggesting an increased need for resuscitation among home births.<sup>2</sup> Therefore, the personnel, training, and equipment available for neonatal resuscitation represent other possible contributors to the excessive neonatal mortality rate among planned home births. Finally, we note that there may well be other unrecognized factors contributing to the higher neonatal death rate among planned home births.

Interestingly, our metaanalysis noted similar perinatal mortality rates by intended delivery site, both overall, as well as among nonanomalous offspring. This result is not surprising considering the low-risk nature of the antecedent pregnancies. However, it is an unexpected finding given the increased neonatal mortality rate observed with planned

home delivery. The apparent discordance may result from the differences in obstetric risk among women planning home vs hospital births. A study published after our analysis found similar perinatal mortality rates in planned home and hospital deliveries. However, adjusting the perinatal mortality ratio for the later gestational ages at delivery and greater birthweights among home births demonstrated higher standardized perinatal mortality ratios among planned home deliveries, particularly among those requiring transfer to hospital.<sup>22</sup> Such an adjustment could not be performed in the current analysis without patient-level data. However, one may speculate that similar findings would be noted based on the later gestational age at birth and greater birthweights seen in our analysis among planned home vs planned hospital births. In contrast, we were able to estimate the population-based attributable risk of neonatal death due to home birth. The absolute risk was small, reflecting the low prevalence of home birth and rarity of the outcome, despite its significantly increased OR.

A paucity of data in the original studies precluded a more in-depth examination

of contributors to the perinatal mortality rates described in this metaanalysis. Potentially valuable insights could result from evaluating antepartum vs intrapartum stillbirths, as well as potentially preventable deaths. Interestingly, 2 Dutch studies observed no relationship between potentially avoidable perinatal deaths and delivery setting (home vs hospital) or birth attendant (midwife vs physician).<sup>23,24</sup> However, a recent Australian study identified an increased rate of intrapartum perinatal deaths among planned home deliveries, one third of which were attributed to asphyxia, contrasting only 3.6% of intrapartum perinatal deaths among planned hospital births.<sup>22</sup>

The maternal mortality rate arguably represents the ultimate measure of child-birth safety. The current study could not perform metaanalysis of maternal mortality by planned delivery location because no deaths were described among studies reporting this outcome. The absence of maternal deaths is not surprising considering the number of deliveries comprising the study populations. Thus, more data are necessary before drawing any conclusions regarding the maternal

mortality rates of planned home and planned hospital delivery.

The current metaanalysis shows that planned home compared to planned hospital births are associated with significantly less maternal and newborn medical intervention and morbidity particularly among selected low-risk women cared for by highly trained and regulated midwives who are integrated into the health care system. These findings are notable in that our analysis by planned delivery site confirms many of the observations of a recent cohort study evaluating outcomes by actual delivery location.<sup>2</sup> At first glance, these results are not surprising for several reasons. Many women choose home birth, at least in part to avoid pharmacologic analgesia and medical technology.<sup>4,25-30</sup> Most women considered to be home birth candidates exhibit low obstetric risk and should therefore anticipate more favorable outcomes than women choosing or requiring a planned hospital delivery. Finally, most home births are attended by midwives, a group demonstrating distinctly different obstetric practice patterns from physicians performing most in-hospital deliveries.<sup>31-34</sup> A systematic review and metaanalysis of randomized trials of midwife-led vs other care models confirms less medical intervention and improved perinatal outcomes in the former group.<sup>35</sup> Importantly, these trials included hospital but not home births.

Women, particularly low-risk parous individuals, choosing home birth are in large part successful in achieving their goal of delivering with less morbidity and medical intervention than experienced during hospital-based childbirth. Of significant concern, these apparent benefits are associated with a near tripling of the neonatal mortality rate among nonanomalous infants. These results confirm and complement those of prior large cohort studies assessing outcomes by actual birth location, suggesting generalizability to and value in counseling low-risk women considering home birth particularly with highly trained, regulated midwives who are fully integrated into existing health care systems. Therefore, these data may be of limited applicability to women opting

for home birth in the United States.<sup>36</sup> The large number of outcomes for which heterogeneity was present suggests that such results should be interpreted with caution. Finally, one must appreciate that the lower obstetric risk characterizing women self-selecting planned home birth likely underestimates the risk and overestimates the benefit of this delivery choice.

Future research needs to be directed at identifying contributors to and reducing the apparently excessive neonatal mortality among planned home births. Data regarding maternal mortality, maternal and newborn readmission rates and indications, and newborn neurologic injury are insufficient for evaluation and comparison. Comprehensive economic analyses by planned birth location are also lacking.<sup>37</sup> Ideally, the results of such work will contribute to an obstetric and newborn best practices model benefiting women and children regardless of chosen birth location. ■

#### REFERENCES

- Martin JA, Hamilton BE, Sutton PD, et al. Births: final data for 2006. *Natl Vital Stat Rep* 57. Available at [www.cdc.gov/nchs/data/nvsr/nvsr57/nvsr57\\_07.pdf](http://www.cdc.gov/nchs/data/nvsr/nvsr57/nvsr57_07.pdf). Accessed June 1, 2010.
- Wax JR, Pinette MG, Cartin A, Blackstone J. Maternal and newborn morbidity by birth facility among selected United States 2006 low-risk births. *Am J Obstet Gynecol* 2010;202:152.e1-5.
- American College of Obstetricians and Gynecologists. ACOG statement of policy: home births in the United States. Available at: [http://www.acog.org/publications/policy\\_statements/sop0705.cfm](http://www.acog.org/publications/policy_statements/sop0705.cfm). Accessed Jan. 27, 2009.
- Ackermann-Lieblich U, Voegeli T, Günter-Witt K, et al. Home versus hospital deliveries: follow up study of matched pairs for procedure and outcome. *BMJ* 1996;313:1313-8.
- Shearer JML. Five year prospective survey of risk of booking for a home birth in Essex. *BMJ* 1985;291:1478-80.
- Wiegars TA, Keirse MJNC, van der Zee J, Berghs GAH. Outcome of planned home and planned hospital births in low risk pregnancies: prospective study in midwifery practices in the Netherlands. *BMJ* 1996;313:1309-13.
- Lindgren HE, Radestad IJ, Christensson K, Hildengsson IM. Outcomes of planned home births compared to hospital births in Sweden between 1992 and 2004: a population-based register study. *Acta Obstet Gynecol* 2008;87:751-9.
- Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology. *JAMA* 2000;283:2008-12.
- Zaza S, Wright-DeAguro LK, Briss PA, et al. Data collection instrument and procedure for systematic reviews in the guide to community preventive services. *Am J Prev Med* 2000;18:44-74.
- Woodcock HC, Read AW, Bower C, Stanley FJ, Moore DJ. A matched cohort study of planned home and hospital births in Western Australia 1981-1987. *Midwifery* 1994;10:125-35.
- Hutton EK, Reitsma AH, Kaufman K. Outcomes associated with planned home and planned hospital births in low-risk women attended by midwives in Ontario, Canada, 2003-2006: a retrospective cohort study. *Birth* 2009;36:180-9.
- Janssen PA, Saxell L, Page LA, Klein MC, Liston RM, Lee SK. Outcomes of planned home birth with registered midwife versus planned hospital birth with midwife or physician. *CMAJ* 2009;181:377-83.
- Koehler NU, Solomon DA, Murphy M. Outcomes of a rural Sonoma county home birth practice: 1976-1982. *Birth* 1984;11:165-9.
- Dowswell T, Thornton JG, Hewison J, Lilford RJL. Should there be a trial of home versus hospital delivery in the United Kingdom? *BMJ* 1996;312:753-7.
- Pang JWY, Heffelfinger JD, Huang GJ, Benedetti TJ, Weiss NJ. Outcomes of planned home births in Washington State: 1989-1996. *Obstet Gynecol* 2002;100:253-9.
- deJong A, van der Goes BY, Ravelli ACJ, et al. Perinatal mortality and morbidity in a nationwide cohort of 529,688 low-risk planned home and hospital births. *BJOG* 2009;116:1177-84.
- Janssen PA, Lee SK, Ryan EM, et al. Outcomes of planned home births versus planned hospital births after regulation of midwifery in British Columbia. *CMAJ* 2002;166:315-23.
- Pasupathy D, Wood AM, Pell JP, Fleming M, Smith GCS. Rates of and factors associated with delivery-related perinatal death among term infants in Scotland. *JAMA* 2009;302:660-8.
- Crotty M, Ramsay AT, Smart R, Chan A. Planned homebirths in South Australia 1976-1987. *Med J Aust* 1990;153:664-71.
- Bastian H, Keirse MJNC, Lancaster PAL. Perinatal death associated with planned home birth in Australia: population based study. *BMJ* 1998;317:384-8.
- Vintzileos AM. Evidence-based compared with reality-based medicine in obstetrics. *Obstet Gynecol* 2009;113:1335-40.
- Kennare RM, Keirse M, Tucker GR, Chan AC. Planned home and hospital births in South Australia, 1991-2006: differences in outcomes. *Med J Aust* 2010;192:76-80.
- Wolleswinkel-van den Bosch JH, Vredevoogd CB, Borkent-Polet M, et al. Substandard factors in perinatal care in the Netherlands: a regional audit of perinatal deaths. *Acta Obstet Gynecol Scand* 2002;81:17-24.
- DeReu PAOM, Nijhuis JG, Oosterbaan HP, Eskes TKAB. Perinatal audit on avoidable mor-

- tality in a Dutch rural region: a retrospective study. *Eur J Obstet Gynecol Reprod Biol* 2000;88:65-9.
- 25.** Hildingsson I, Waldenstrom U, Radestad I. Swedish women's interest in home birth and in-hospital birth center care. *Birth* 2003;30:11-22.
- 26.** vanderHulst LAM, vanTeijlingen ER, Bonsel GJ, Eskes M, Bleker OP. Does a pregnant woman's intended place of birth influence her attitudes toward and occurrence of obstetric interventions? *Birth* 2004;31:28-33.
- 27.** Ogden J, Shaw A, Zander L. Women's memories of home-birth 3-5 years on. *Br J Midwifery* 1997;5:208-11.
- 28.** Neuhaus W, Piroth C, Kiencke P, Göhring U, Mallman P. A psychosocial analysis of women planning birth outside hospital. *J Obstet Gynaecol* 2002;22:143-9.
- 29.** Abernathy TJ, Lentjes DM. Planned and unplanned home births and hospital births in Calgary, Alberta, 1984-87. *Public Health Rep* 1989;104:373-7.
- 30.** Boucher D, Bennett C, McFarlin B, Freeze R. Staying home to give birth: why women in the United States choose home birth. *J Midwifery Womens Health* 2009;54:119-26.
- 31.** Fraser W, Hatem-Asmar M, Krauss I, et al. Comparison of midwifery care to medical care in hospitals in the Quebec pilot study: clinical indicators. *Can J Public Health* 2000;91:15-11.
- 32.** Rosenblatt R, Dobie SA, Hart CG, et al. Interspecialty differences in the obstetric care of low-risk women. *Am J Public Health* 1997;87:344-51.
- 33.** Reime B, Klein MC, Kelly A, et al. Do maternity care provider groups have different attitudes toward birth? *BJOG* 2004;111:1388-99.
- 34.** Janssen PA, Ryan EM, Etches DJ, Klein MC, Reime B. Outcomes of planned hospital birth attended by midwives compared with physicians in British Columbia. *Birth* 2007;34:140-7.
- 35.** Hatem M, Sandall J, Devane D, Soltani H, Gates S. Midwife-led versus other models of care for childbearing women. *Cochrane Database Syst Rev* 2008;4:CD004667.
- 36.** Wax JR, Pinette MG, Cartin A. Home versus hospital birth: process and outcome. *Obstet Gynecol Surv* 2010;65:132-40.
- 37.** Henderson J, Petrou S. Economic implications of home births and birth centers: a structured review. *Birth* 2008;35:136-46.